

FUNCTIONAL CHARACTERISATION OF A
NOVEL IMMUNE MODULATORY
MOLECULE FROM *FASCIOLA HEPATICA*

by

Raquel Alvarado B.Sc. (Hons)

A Thesis Submitted for the Degree of

Doctor of Philosophy in Science

School of Medical and Molecular Biosciences, Faculty of Science,

University of Technology Sydney, Australia.

2014

CERTIFICATE OF AUTHORSHIP/ORIGINALITY

I certify that the work in this thesis has not previously been submitted for a degree nor has it been submitted as part of requirements for a degree except as fully acknowledged within the text.

I also certify that the thesis has been written by me. Any help that I have received in my research work and the preparation of the thesis itself has been acknowledged. In addition, I certify that all information sources and literature used are indicated in the thesis.

Signature of Student: 

Date: 22nd December, 2014

ACKNOWLEDGEMENTS

Science has fascinated me ever since I was a child, therefore I am very grateful to God for not only giving me the opportunity, but also the willingness and strength to pursue my studies in this field that I truly enjoy. I consider myself very fortunate for being able to do this PhD project, which despite being my greatest challenge as a student, has rewarded me with getting to know so many people who have generously shared their knowledge and friendship with me.

To my parents, I thank them for all their love, for listening, encouraging me and always giving me their complete support and to my mum in particular for being my best friend. Without them I would simply not have been able to achieve this goal.

I thank Bronwyn O'Brien, Sheila Donnelly and Mark Robinson, for giving me the privilege of working with them. It was Bronwyn's enthusiastic Immunology lectures that first showed me how interesting this field can be. I really appreciate the direct involvement in the laboratory of Sheila and Mark, who patiently dedicated a lot of their time to teach me new techniques and help me with experiments. I want to thank the three of them for giving me such an interesting project, for teaching me so much, checking my drafts, giving their valuable feedback, for all their guidance and overall for being such excellent supervisors. I'm also very grateful to Joyce To for helping me so much in this project and for teaching me her clever modifications of experimental techniques. To Maria Lund, Andrew Hutchinson and Paddy McCauley-Winter, I thank all their assistance in flow cytometry and their valuable advice. I am very proud to have been part of this research team, whose members were so kind to me and I wish them all very successful futures with the hope that our friendship endures a very long time.

In these years I have also met fellow postgraduate students with whom I have acquired what I would love to be lifelong friendships, and with whom I have shared unforgettable moments. I want to thank them for being so wonderful, especially to Maria Lund, Alex Gale and Rita Rapa for letting me count on them in good and difficult moments.

I would also like to express my gratitude to the UTS technical staff, especially Mike Johnson and Lynne Turnbull for their assistance in microscopy. Also, I thank Ashley Mansell, Anita Pinar and Suat Dervish for their contribution to my project and the science faculty for granting me a postgraduate research scholarship.

PUBLICATIONS ASSOCIATED WITH THIS THESIS

Journal Publications

Mark W. Robinson, **Raquel Alvarado**, Joyce To, Andrew Hutchinson, Stephanie N. Dowdell, Maria Lund, Lynne Turnbull, Cynthia B. Witchurch, Bronwyn A. O'Brien, John P. Dalton and Sheila Donnelly, 2012; A helminth cathelicidin-like protein suppresses antigen processing and presentation in macrophages via inhibition of lysosomal vATPase. *FASEB Journal*. 26, 4614-4627.

Raquel Alvarado, Bronwyn O'Brien, Akane Tanaka, John P. Dalton, Sheila Donnelly, 2014; A parasitic helminth-derived peptide that targets the macrophage lysosome is a novel therapeutic option for autoimmune disease. *Immunobiology*. (In press, DOI 10.1016/j.imbio.2014.11.008).

Raquel Alvarado, Joyce To, Anita Pinar, Ashley Mansell, Mark Robinson, Bronwyn O'Brien, John Dalton, Sheila Donnelly, 2014; The *F. hepatica* cathelicidin like peptide (FhHDM-1) modulates the activation of the NLRP3 inflammasome in macrophages. (Manuscript in preparation).

Conference presentations

Raquel Alvarado, Maria Lund, Mark W. Robinson, Andrew Hutchinson, Joyce To, John P. Dalton, Bronwyn O'Brien and Sheila Donnelly, 2012; Presentation entitled: A novel molecule secreted by *Fasciola hepatica* modulates the activity of innate immune cells. 42nd Annual Scientific Meeting of the Australasian Society for Immunology.

Raquel Alvarado, Joyce To, Maria Lund, Mark W. Robinson, Andrew Hutchinson, Bronwyn O'Brien, John P. Dalton and Sheila Donnelly, 2013; Presentation entitled: A novel molecule secreted by the parasite *F. hepatica* modulates the response of macrophage NLRP3 inflammasome. 27th Annual Conference of the European Macrophage & Dendritic Cell Society.

TABLE OF CONTENTS

Certificate of authorship/originality	i
Acknowledgements	ii
Publications associated with this thesis.....	iii
Table of contents	iv
List of illustrations	ix
List of tables.....	xi
Abbreviations	xii
Abstract	xvi
Chapter 1 General Introduction.....	1
1.1. Versatility of the mammalian immune system.....	1
1.2. Helminth parasites are master regulators of the mammalian immune system	2
1.3. Characterisation of immune modulatory components of helminth excretory/secretory products	5
1.3.1. Proteases.....	6
1.3.2. Protease inhibitors	7
1.3.3. IPSE/alpha-1, Omega-1 and other helminth glycans	9
1.3.4. Cytokine homologues.....	10
1.3.5. Antioxidants	11
1.3.6. ES-62 a phosphorylcholine (PC) containing glycoprotein.....	12
1.3.7. Additional ES products that are immune modulators	13
1.4. <i>Fasciola hepatica</i> : A model of helminth-induced immune-modulation.....	18
1.4.1. FhCL1.....	20
1.4.2. FhPrx	21

1.4.3. FhHDM-1	22
1.5. FhHDM-1 is a helminth-derived peptide with homology to the mammalian cathelicidin defence peptides	24
1.5.1. A proposed mechanism of action for FhHDM-1	27
Chapter 2 General Materials & Methods	29
2.1. Production of RecFhHDM-1, sFhHDM-1 and anti-FhHDM-1 antibody	30
2.2. Cell culture	31
2.2.1. Sterility	31
2.2.2. Cell lines.....	31
2.2.2.1. RAW264.7 macrophages	31
2.2.2.2. ASC macrophages	31
2.2.3. Primary cells.....	32
2.2.3.1. Murine Bone Marrow-Derived Macrophages (BMDMs)	32
2.2.3.1.1. Isolation of bone marrow cells	32
2.2.3.1.2. Differentiation of Bone Marrow-Derived Macrophages	32
2.2.3.2. Human Monocyte Isolation and Macrophage Differentiation	32
2.3. Immunofluorescence Confocal Microscopy	33
2.3.1. NPG Antifade Mounting Media preparation.....	35
2.4. Flow Cytometry.....	35
2.5. ELISAs	35
2.6. Statistical Analysis	36
Chapter 3 Characterisation of the interaction between FhHDM-1 and macrophages	37
3.1. Introduction	37
3.2. Specific methods	39
3.2.1. Cholesterol binding assay.....	39
3.2.2. Immunofluorescent Confocal Microscopy	39
3.2.2.1. Co-localisation of FhHDM-1 with lipid rafts.....	39

3.2.2.2. Inhibitor Studies	40
3.2.2.3. Co-localisation of FhHDM-1 with organelle markers	40
3.2.2.3.1. Localisation of FhHDM-1 with caveolae, Golgi and mitochondria	40
3.2.2.3.2. Temporal localisation of FhHDM-1 with early endosomes.....	41
3.2.2.3.3. Temporal localisation of FhHDM-1 with late endosomes/lysosomes	41
3.3. Results	43
3.3.1. FhHDM-1 binds to cholesterol and interacts with lipid rafts.....	43
3.3.2. FhHDM-1 is actively endocytosed by a cytoskeletal-dependent mechanism.....	45
3.3.3. Endocytosis of FhHDM-1 involves early endosomal and lysosomal co- localisation	50
3.4. Discussion	58
Chapter 4 FhHDM-1 modulates the processing of antigens by macrophages	62
4.1. Introduction	62
4.2. Specific methods	65
4.2.1. Endocytosis and vesicular acidification studies	65
4.2.1.1. Dextran endocytosis studies.....	65
4.2.1.1.1. Flow Cytometry	65
4.2.1.1.2. Confocal microscopy	66
4.2.2. Effects of FhHDM-1 on antigen processing and presentation	66
4.2.2.1. Effects of sFhHDM-1 on DQ Ovalbumin processing.....	66
4.2.2.2. Effects of FhHDM-1 on antigen presentation to transgenic murine cells.....	67
4.2.2.2.1. Antigen processing studies.....	67
4.2.2.2.2. T cell isolation.....	67
4.2.2.2.3. Antigen presentation studies	68
4.2.3. Effects of FhHDM-1 on MHCII surface expression by BMDMs.....	68
4.3. Results	69

4.3.1. FhHDM-1 reduces vesicular acidification and enhances endocytosis by macrophages	69
4.3.2. FhHDM-1 reduces antigen processing by macrophages	76
4.3.3. Effects of FhHDM-1 on antigen presentation by macrophages	78
4.4. Discussion	82
Chapter 5 FhHDM-1 reduces NLRP3 inflammasome activation in macrophages	87
5.1. Introduction	87
5.2. Specific methods	91
5.2.1. FhHDM-1 effects on NLRP3 Inflammasome Activation	91
5.2.1.1. NLRP3 activation	91
5.2.1.2. Cytokine detection by ELISA	91
5.2.1.3. Detection of activated IL-1 β and Caspase-1 by Western Blot	91
5.2.1.3.1. TCA supernatant protein precipitation	92
5.2.1.3.2. Gel Electrophoresis	92
5.2.1.3.3. Western Blotting	92
5.2.2. ASC speck formation	93
5.2.3. FhHDM-1 effects on lysosomal integrity	93
5.2.3.1. Lysosomal stability	93
5.2.3.1.1. DQ Ova compartmentalisation	93
5.2.3.2. Cathepsin B activity	94
5.3. Results	95
5.3.1. FhHDM-1 inhibits lysosomal-dependent NLRP3 activation	95
5.3.1.1. FhHDM-1 reduces NLRP3 inflammasome activation by ALUM	95
5.3.1.2. FhHDM-1 reduces NLRP3 inflammasome activation by Nano-SiO ₂	96
5.3.2. FhHDM-1 prevents NLRP3 dependent ASC oligomerisation	98
5.3.3. Effects of FhHDM-1 on lysosomal integrity and stability after ALUM-induced inflammasome activation	101

5.3.4. Effects of FhHDM-1 on LPS priming.....	105
5.4. Discussion	107
Chapter 6 General Discussion.....	111
References	114

LIST OF ILLUSTRATIONS

Figure 1.1 Characterization of FhHDM-1 and structural homology to LL-37.	23
Figure 3.1 FhHDM-1 binds to cholesterol and co-localises with lipid rafts in the plasma membranes of macrophages.	44
Figure 3.2 FhHDM-1 is internalised by macrophages and localises in close proximity to cytoskeletal networks.	46
Figure 3.3 Internalisation of FhHDM-1 by macrophages is an active process dependent upon cholesterol, as well as actin and microtubule networks.	49
Figure 3.4 FhHDM-1 endocytosis was not mediated by caveolae.	51
Figure 3.5 FhHDM-1 co-localised with early endosomes.	52
Figure 3.6 FhHDM-1 co-localised with endolysosomes of fixed RAW264.7 macrophages.	54
Figure 3.7 FhHDM-1 co-localised with endolysosomes of live BALBc BMDMs.	55
Figure 3.8 FhHDM-1 did not co-localise with the Golgi apparatus or mitochondria in RAW264.7 macrophages.	57
Figure 4.1 Macrophage pre-treatment with FhHDM-1 decreased dextran detection in acidified vesicles but not its endocytosis.	72
Figure 4.2 Simultaneous incubation of macrophages with FhHDM-1 did not affect the detection of dextran in acidified vesicles, but it did increase the uptake of dextran.	73
Figure 4.3 Macrophages treated with FhHDM-1 can endocytose dextran and simultaneous incubation with both molecules enhanced the co-localisation of FhHDM-1 and dextran.	75
Figure 4.4 Simultaneous incubation with FhHDM-1 and ovalbumin reduces the ability of macrophages to process antigen.	77
Figure 4.5 FhHDM-1 reduces the ability of macrophages to process antigenic peptides for their presentation to T cells.	79
Figure 4.6 FhHDM-1 did not alter MHCII expression levels by macrophages.	81
Figure 5.1 NLRP3 activation pathway.	89

Figure 5.2 FhHDM-1 reduced NLRP3 inflammasome activation induced by lysosomal destabilising agents.	97
Figure 5.3 FhHDM-1 reduced ASC speck formation.	100
Figure 5.4 FhHDM-1 does not prevent the lysosomal destabilisation induced by ALUM.	103
Figure 5.5 FhHDM-1 treatment of macrophages reduced cathepsin B activity.	104
Figure 5.6 FhHDM-1 reduced LPS priming efficiency in macrophages.	106

LIST OF TABLES

Table 1.1 Classification of identified helminth ES components and their immune modulatory/evasive effects.	16
Table 2. 1 General materials and reagents.	29
Table 2.2 List of antibodies and dyes used for confocal microscopy experiments.....	34

ABBREVIATIONS

ACEC	Animal Care and Ethic Committee
ADCC	Antibody dependent cell mediated cytotoxicity
AIM2	Absent in melanoma 2
ALUM	Aluminium salts
AMPs	Antimicrobial peptides
APC(s)	Antigen presenting cell(s)
ARC	Animal Resources Centre
ASC	Apoptosis associated speck like protein
ATCC	American Type Culture Collection
Az	Azide
BcR	B cell receptor
BMDMs	Bone marrow derived macrophages
BSA	Bovine serum albumin
CAT	Catalases
Cav-1	Caveolin-1
ChTx	Cholera toxin subunit B
CLIC	Clathrin-independent non-caveolar pathway
CO₂	Carbon dioxide
DAMPs	Damage-associated molecular patterns
DAPI	4'6-diamidino-2 phenylindole, dilactate
DMSO	Dimethyl sulfoxide
DC(s)	Dendritic cell(s)
<i>E. coli</i>	<i>Escherichia coli</i>
ELISA	Enzyme-linked immunosorbent assay
ES	Excretory secretory products
FACS	Fluorescence-activated cell sorting (flow cytometry)
FBS	Foetal bovine serum
FhCL1	<i>Fasciola hepatica</i> cathepsin L-1
FhES	<i>Fasciola hepatica</i> excretory/secretory products
FhHDM-1	<i>Fasciola hepatica</i> helminth defence molecule-1
FhHDM-1p2	FhHDM-1 peptide 2
FhPrx	<i>Fasciola hepatica</i> peroxiredoxin

Geo Mean	Geometric mean
GPx	Glutathione peroxidase
hCAP18	Human cationic antimicrobial protein 18kDa
HDMs	Helminth defence molecules
HDPs	Host defence peptides
HEPES	4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid
HVS	<i>Herpesvirus saimiri</i>
IAPP	Islet amyloid polypeptide
IFNγ	Interferon gamma
Ig	Immunoglobulin
IL	Interleukin
IMDM	Iscove's modified Dulbecco's medium
IPSE	IL-4 inducing principle of schistosome eggs
kDa	Kilo daltons
LBP	LPS-binding protein
LNFP III	Lacto-N-fucopentaose III
LPS	Lipopolysaccharide
MAPKs	Mitogen-activated protein kinases
MCD	Methyl- β -cyclodextrin
M-CSF	Macrophage colony stimulating factor
MHC	Major histocompatibility complex
MIC	Minimal concentration capable of inhibiting visible microbial growth
MIF	Migration inhibitory factor
MPR	Mannose phosphate receptor
mRNA	Messenger ribonucleic acid
MS	Multiple sclerosis
MSU	Mono sodium urate
MW	Molecular weight
MyD88	Myeloid differentiation factor 88
N/A	Not applicable
Na₂CO₃	Sodium carbonate
Nano-SiO₂	Silicon dioxide (nanoparticles)
NH₄Cl	Ammonium chloride
NPG	N-propyl gallate microscopy mounting media

NLRs	NACHT-leucine-rich repeat receptors
(NOD)-like receptors	Nucleotide-binding oligomerisation domain protein like receptors
OD	Optical density
O/N	Overnight
OPep	Ovalbumin peptide
Ova	Ovalbumin
PAMPS	Pathogen-associated molecular patterns
PBMCs	Peripheral blood mononuclear cells
PBS	Phosphate buffered saline
PC	Phosphorylcholine
PFA	Paraformaldehyde
PGE₂	Prostaglandin E2
PI	Peak I
PII	Peak II
PI 3-K	Phosphoinositide 3 kinase
Prx	Peroxiredoxin
RecFhHDM-1	Recombinant <i>Fasciola hepatica</i> helminth defence molecule 1
RELM-α	Resistin-like molecule-alpha
RIPA	Radioimmunoprecipitation Assay Buffer
ROS	Reactive oxygen species
RPMI	Roswell Park Memorial Institute 1640 medium
RP-HPLC	Reversed-phase high performance liquid chromatography
RT	Room temperature
SmCB1	<i>Schistosoma mansoni</i> cathepsin B
SDS-PAGE	Sodium dodecyl sulphate- polyacrylamide gel electrophoresis
SEA	Soluble egg antigens
SEMs	Standard errors of the means
sFhHDM-1	Synthetic <i>Fasciola hepatica</i> helminth defence molecule-1
SiO₂	Silicon dioxide (nanoparticles)
SOD	Superoxide dismutases
T1D	Type I diabetes
TBS	Tris buffer saline
TcR	T cell receptor
TGF-β	Transforming growth factor beta

TGN	Trans-Golgi network
Th	T helper cells
Tip	Tyrosine kinase interacting protein
TLR	Toll like receptor
TNF	Tumour necrosis factor
TMB	3,3',5,5'-Tetramethylbenzidine liquid substrate system for ELISA
Treg	T regulatory cells
TRIF	TIR domain-containing adaptor inducing IFN- β
TX100	Triton X 100
vATPase	Vacuolar adenosine triphosphatase
v/v	Volume / volume
w/v	Weight / volume

ABSTRACT

The ability of tissue dwelling helminth parasites to induce chronic long term infections, is enabled by the establishment of T helper 2/ regulatory T cell (Th2/Treg) immune responses within their mammalian hosts. Such responses prevent the expulsion of the parasites, whilst simultaneously avoiding excessive inflammation/fibrosis arising within the host, as a consequence of tissue damage induced by helminth migration. Importantly, helminths excrete and secrete a series of molecules (collectively known as ES products), which not only play major roles in parasite biology, but also exert direct immune modulatory functions, promoting the establishment of Th2/Treg immunity. The trematode, *Fasciola hepatica*, is an excellent model of helminth-mediated immune modulation, because it induces a very rapid switch towards Th2 responses in its mammalian hosts and inhibits Th1 immunity. Fractionation of the ES products of *F. hepatica* has identified three major immune modulatory components: the protease cathepsin L1, the antioxidant peroxiredoxin, and a previously uncharacterised peptide, FhHDM-1.

Structural analysis of FhHDM-1 revealed a close resemblance to the cathelicidin, LL-37, a well characterised mammalian immune-modulating peptide. Therefore, a putative immune modulatory role for FhHDM-1 was explored in this project. Immunofluorescent confocal microscopy demonstrated that FhHDM-1 interacted with macrophage lipid rafts, prior to being actively internalised by cholesterol- and cytoskeletal network-dependent endocytosis, with progressive compartmentalisation of the peptide into early endosomes and endolysosomal vesicles. Flow cytometry studies indicated that, once internalised, FhHDM-1 enhanced the rate of endocytosis of dextran by macrophages. Despite this, FhHDM-1 was found to impair the acidification of macrophage endolysosomes and as a consequence, the efficient processing and subsequent presentation of ovalbumin to T cells was prevented, as assessed by decreased detection of digested fluorescent ovalbumin and reduced IL-2 secretion by transgenic CD4⁺ T cells. Additionally, FhHDM-1 impaired NLRP3 inflammasome activation by lysosomal disruptive agents in macrophages. This was found to be a consequence of reduced cathepsin B activity (due to FhHDM-1 induced suboptimal lysosomal acidification), which was incapable of stimulating inflammasome complex formation, thus avoiding IL-1 β and caspase-1 cleavage.

These findings suggest that by targeting endolysosomal activity, FhHDM-1 limits macrophage function. Therefore, the current study is the first to demonstrate that FhHDM-1 possesses immune modulatory properties, which are directed by a mechanism not previously described for a helminth-secreted cathelicidin-like peptide.